

# Molecular mechanisms of Cl<sup>-</sup> transport in fishes: New insights and their evolutionary context

Ciaran A. Shaughnessy<sup>1</sup>  | Jason P. Breves<sup>2</sup> 

<sup>1</sup>Department of Pediatrics, National Jewish Health, Denver, Colorado, USA

<sup>2</sup>Department of Biology, Skidmore College, Saratoga Springs, New York, USA

## Correspondence

Jason P. Breves, PhD, Department of Biology, Skidmore College, 815 N. Broadway, Saratoga Springs, NY 12866, USA.

Email: [jbreves@skidmore.edu](mailto:jbreves@skidmore.edu)

## Funding information

Eugene F. and Easton M. Crawford Charitable Lead Unitrust to National Jewish Health; National Science Foundation, Division of Integrative Organismal Systems, Grant/Award Number: IOS-1755131

## Abstract

With remarkably few exceptions, aquatic vertebrates maintain internal Cl<sup>-</sup> homeostasis despite strong and sometimes fluctuating Cl<sup>-</sup> concentration gradients between extracellular fluids and external environments. In this “Perspective,” we discuss recent advances in the understanding of epithelial Cl<sup>-</sup> transport at the molecular level within key osmoregulatory organs in fishes. New insights into mechanisms for epithelial Cl<sup>-</sup> transport in basal lineages are highlighted to provide an evolutionary context. We describe Cl<sup>-</sup> transport processes that employ: cystic fibrosis transmembrane conductance regulator, cation-chloride cotransporters, voltage-gated chloride channels, and chloride-anion exchangers. As the collective understanding of Cl<sup>-</sup> transport processes continues to expand, investigators are equipped to more precisely characterize how endocrine factors promote hydromineral balance. We, therefore, conclude our discussion by paying special attention to recently defined roles for prolactin and corticosteroids in the regulation of Cl<sup>-</sup> transport in basal and derived clades.

## KEYWORDS


gill, hormones, ionocytes, lamprey, sturgeon, teleost

## 1 | INTRODUCTION

Fishes are grouped into three classes: Agnatha (jawless fishes), Chondrichthyes (cartilaginous fishes), and Osteichthyes (bony fishes) (Figure 1). Teleosts (class Osteichthyes; subclass Actinopterygii; infra-class Neopterygii; division Teleostei) and lampreys (members of Class Agnatha) typically maintain extracellular fluids between 270 and 400 mOsm/kg, with the major dissolved ions, Na<sup>+</sup> and Cl<sup>-</sup>, maintained between approximately 130–180 and 100–150 mmol/l, respectively (Ferreira-Martins et al., 2016; Marshall & Grosell, 2006). Freshwater (FW) and marine environments offer formidable challenges to the maintenance of ionic and osmotic homeostasis. Teleosts and lampreys residing in FW are hyperosmotic to the dilute external environment; therefore, they are dually threatened by diffusive ion losses and excessive hydration. To maintain hydromineral balance, the gills absorb Na<sup>+</sup> and Cl<sup>-</sup> against strong concentration gradients while the kidney reabsorbs ions and exhibits high glomerular filtration rates to produce large volumes of dilute urine (Marshall &

Grosell, 2006). Teleosts and lampreys residing in seawater (SW), on the other hand, are hyposmotic to the external environment and face the passive gain of ions and dehydration. In turn, they must drink ambient SW that is initially desalinated by the esophagus to enable subsequent solute-linked water absorption by the intestine (Barany et al., 2020; Grosell, 2006; Takei et al., 2017). The gills are specialized for Na<sup>+</sup> and Cl<sup>-</sup> secretion while the kidney exhibits low glomerular filtration rates and produces small volumes of isosmotic urine (Marshall & Grosell, 2006). The branchial, gastrointestinal, and renal epithelia of both FW- and SW-acclimated fishes must leverage various pathways to actively transport Cl<sup>-</sup>, the major anion in extracellular fluids, to sustain hydromineral balance of the organism.

In the past two decades, a rapid increase in the genomic resources available for the study of teleost fishes has illuminated the diversity of molecular strategies for osmoregulation. Recent sequencing efforts for pirarucu (*Arapaima gigas*; Vialle et al., 2018), spotted gar (*Lepisosteus oculatus*; Braasch et al., 2016), sterlet sturgeon (*Acipenser ruthenus*; Cheng et al., 2019; Du et al., 2020), bichir (*Polypterus senegalus*;



Chiu et al., 2004), African coelacanth (*Latimeria chalumna*; Amemiya et al., 2013), inshore hagfish (*Eptatretus burgeri*; Yamaguchi et al., 2020), and sea lamprey (*Petromyzon marinus*; Smith et al., 2013, 2018) will now support future investigations into the mechanisms underlying  $\text{Cl}^-$  transport in basal actinopterygian and pre-actinopterygian lineages. Here, we first present recent insights into the molecular mechanisms underlying  $\text{Cl}^-$  transport in fishes with special reference to their operation in basal lineages. We then discuss how these  $\text{Cl}^-$  transport pathways are functionally controlled by the pituitary hormone, prolactin, and corticosteroids.

## 2 | MOLECULAR MECHANISMS OF $\text{Cl}^-$ TRANSPORT IN FISHES


### 2.1 | Cystic fibrosis transmembrane conductance regulator (Cftr)

For decades it has been known that marine/SW-acclimated teleosts express Cftr on the apical surface of branchial ionocytes where it serves as the conduit for active  $\text{Cl}^-$  secretion (Marshall & Singer, 2002). In these same ion-secreting cells, termed “SW-type” ionocytes,  $\text{Na}^+/\text{K}^+$ -ATPase (Nka) and  $\text{Na}^+/\text{K}^+/\text{2Cl}^-$  cotransporter 1 (Nkcc1) are coexpressed in the basolateral membrane to energize and facilitate the  $\text{Na}^+$ - and  $\text{K}^+$ -coupled passage of  $\text{Cl}^-$  from blood plasma into ionocytes (Figure 2). The coordinated operation of Cftr, Nkcc1, and Nka is represented in multiple  $\text{Cl}^-$ -secreting vertebrate epithelia, including shark rectal gland, teleost gill, avian/reptilian salt glands, and the mammalian intestinal crypt and airway. Recent characterizations of functionally divergent Cftr isoforms in fishes have provided new insight into the structural and functional evolution of Cftr in vertebrates.

Given its rich population of ionocytes and structural advantages for electrophysiological analyses, the opercular membrane of

mummichog (*Fundulus heteroclitus*) has a long-standing record as an in vitro model for the study of  $\text{Cl}^-$  transport (Karnaky et al., 1977; Marshall & Bellamy, 2010; Marshall & Singer, 2002). Fittingly, the first teleost Cftr was described in the opercular epithelium, gill, and intestine of SW-acclimated mummichog (Singer et al., 1998). Two genes encoding distinct Cftr isoforms, denoted *cftr1* and *cftr2*, were subsequently discovered in Atlantic salmon (*Salmo salar*; Chen et al., 2001). With the SW-adaptive role for Cftr1 in teleost gill firmly established, recent investigations focused on discerning the function(s) of Cftr2 in the intestine. Salmon *cftr2* decreased in the intestine during smoltification and SW exposure, patterns indicative of a FW-adaptive role (or a deleterious activity in SW) for this particular isoform (Sundh et al., 2014). In euryhaline sea bream (*Sparus aurata*), intestinal *cftr* expression was similarly diminished by high salinities (Gregório et al., 2013). In Japanese eel (*Anguilla japonica*) where duplicate *cftr* isoforms were also identified, branchial *cftrb* (*cftr1*) was upregulated in SW, whereas intestinal *cftra* (*cftr2*) was diminished in SW (Wong et al., 2016). Given the apical localization of Cftr in mummichog enterocytes, Marshall et al. (2002) proposed that intestinal Cftr supports salt and fluid secretion into the lumen to facilitate the clearance of toxic bacterial flora. This hypothesized role for intestinal Cftr makes intuitive sense; whereas intestinal Cftr activity may be necessary for FW-acclimated teleosts that are not ingesting fluids, it would be unnecessary in the intestine of SW-acclimated teleosts where imbibed fluid is present to facilitate solute-linked water absorption. Functional studies are required to more fully resolve the significance of intestinal Cftr2 expression in FW-acclimated fishes.

In basal fishes, the molecular pathways of branchial  $\text{Cl}^-$  secretion are far less resolved than in teleosts. Our preliminary analysis using BLAST (Altschul et al., 1990) revealed that *cftr* orthologues are present in the genomes of sturgeon, bichir, coelacanth, and hagfish, yet none of these orthologues have been functionally characterized.



**FIGURE 2** Cellular and molecular mechanisms of  $\text{Cl}^-$  transport in key ionoregulatory organs in fishes. Schematic diagrams present molecular mechanisms for  $\text{Cl}^-$  transport that operate in SW and FW gill, FW renal distal tubule, and SW intestine of lampreys, elasmobranchs, sturgeons, and teleosts. Apical and basolateral sides are presented at the top and bottom of cells, respectively. Cell models were derived from protein and/or gene transcript expression patterns depending on data availability. The figure provides a summary of the recently characterized  $\text{Cl}^-$  transport pathways discussed in this “Perspective” and does not intend to serve as an exhaustive depiction of all established or putative pathways. Where  $\text{Cl}^-$  transport is indicated with an arrow and question mark, the presence of a  $\text{Cl}^-$  transport pathway is presumed to exist based on functional observations, but the particular molecular pathway has not been identified. See text for citations. ap, apical; bl, basolateral; Cftr, cystic fibrosis transmembrane conductance regulator; Clc, voltage-gated  $\text{Cl}^-$  channel; FW, freshwater; Ncc,  $\text{Na}^+/\text{Cl}^-$  cotransporter; Nka,  $\text{Na}^+/\text{K}^+$ -ATPase; Nkcc,  $\text{Na}^+/\text{K}^+/2\text{Cl}^-$  cotransporter; Slc26, anion exchanger; SW, seawater; Vha,  $\text{V-H}^+$ -ATPase

A single *cftr* orthologue was previously identified in a genome-wide scan of the sea lamprey, the most basal extant osmoregulating vertebrate (Ren et al., 2015). Ensuing RNAseq analyses in sea lamprey detected low levels of *cftr* expression in all larval, juvenile, and adult tissues except for the intestine of FW-acclimated larvae (Ren et al., 2015). A more recent study confirmed high Cftr expression in the distal intestine of FW-acclimated larvae, and importantly, reported major functional differences between the lamprey Cftr and orthologues from jawed vertebrates. For example, lamprey Cftr exhibits limited  $\text{Cl}^-$  conductance and reduced activation by cAMP versus human Cftr (Cui et al., 2019). Moreover, eel Cftrb and lamprey Cftr exhibit structural divergence in their regulatory domains compared to mammalian Cftrs (Cui et al., 2019; Wong et al., 2016). Intestinal Cftr isoforms of salmon and eel, and the Cftr orthologue of sea lamprey, all possess a reduced number of phosphorylation sites and

reduced responsiveness to protein kinase A (PKA). Thus, PKA-activated  $\text{Cl}^-$  secretion via Cftr does not appear to be a common trait among all vertebrate Cftr isoforms.

Although Cftr orthologues appear to be present throughout the vertebrate phylogeny, it is plausible that its functionality as a  $\text{Cl}^-$  channel has changed throughout vertebrate evolution. Given the limited  $\text{Cl}^-$  conductance of lamprey Cftr and the lack of a *cftr* transcriptional response to SW exposure in the gill (Shaughnessy et al., unpublished), it stands entirely unresolved whether Cftr mediates branchial  $\text{Cl}^-$  secretion at the molecular level in lamprey (Figure 2). Given that SW-acclimated lampreys do in fact utilize Nka- and Nkcc1-expressing SW-type branchial ionocytes (see below), the possibility that an alternate  $\text{Cl}^-$  channel operates as the primary molecular route for branchial  $\text{Cl}^-$  secretion is an exciting avenue for exploration.

## 2.2 | Cation-chloride cotransporters (Slc12 family)

The Slc12 family of electroneutral cation-chloride cotransporters includes  $\text{Na}^+/\text{K}^+/\text{2Cl}^-$  cotransporters (Nkcc1/Slc12a2 and Nkcc2/Slc12a1),  $\text{Na}^+/\text{Cl}^-$  cotransporters (Ncc1/Slc12a3 and Ncc2/Slc12a10), and  $\text{K}^+/\text{Cl}^-$  cotransporters (Kcc1-3/Slc12a4-6). All Slc12 family members are responsible for transporting  $\text{Cl}^-$ , with  $\text{Na}^+$  and/or  $\text{K}^+$ , into a given cell, although they may differ in their transport stoichiometry and cellular location (apical vs. basolateral). The participation of a basolateral Nkcc1 in  $\text{Cl}^-$ -secreting ionocytes of the teleost gill and elasmobranch rectal gland is firmly established (Gagnon et al., 2002; Marshall & Grosell, 2006; Pelis et al., 2001). Nkcc2 and Ncc are expressed within the apical membrane of  $\text{Cl}^-$ -absorptive cells within the gill (Ncc2) and kidney/urinary bladder (Ncc1 and Nkcc2) of FW-acclimated teleosts and within the intestine (Nkcc2) of SW-acclimated teleosts (Figure 2; Kato et al., 2011; Li et al., 2014).

The involvement of Nkcc1 in branchial  $\text{Cl}^-$  secretion by basal fishes has only recently been investigated. Nkcc1 and Nka were increased in sturgeons exposed to salinities  $\geq 25\%$  (McCormick et al., 2020; Sardella & Kültz, 2009; Shaughnessy et al., 2015). In sea lamprey, gill *nkcc1* increased during metamorphosis (the preparatory phase preceding SW migration) and following exposure to SW. Sea lamprey Nkcc1 is colocalized with Nka in presumptive  $\text{Cl}^-$ -secreting branchial ionocytes and functionally critical to maintaining  $\text{Cl}^-$  homeostasis in SW (Shaughnessy & McCormick, 2020).

In SW-acclimated/marine teleosts, the intestine mediates solute-linked water uptake via a suite of ion transporters, channels, and pumps (Grosell, 2006; Sundell & Sundh, 2012). Apically located Nkcc2 mediates the entry of  $\text{Na}^+$  and  $\text{Cl}^-$  into the interior of enterocytes before their subsequent transport across the basolateral membrane (Grosell, 2014; Whittamore, 2012). In Mozambique tilapia (*Oreochromis mossambicus*), Nkcc2 is highly expressed on the luminal surface of the intestine in animals inhabiting SW (Li et al., 2014). In Atlantic salmon, intestinal *nkcc2* increases during smolting and SW acclimation (Sundh et al., 2014). Functional studies exploiting the bumetanide-sensitivity of Nkcc2 have associated *nkcc2* expression with bumetanide-inhibitable current across the intestine of SW-acclimated Japanese eel (Ando & Takei, 2015). In green sturgeon (*A. medirostris*), imbibed  $\text{Cl}^-$  is progressively removed from the gastrointestinal lumen, but the molecular mechanisms for desalination in the sturgeon gut remain unknown (Allen et al., 2009). It was suggested that sea lamprey intestine lacks bumetanide sensitivity (Shaughnessy & McCormick, 2020), and that Nkcc2 might not be a major pathway for intestinal  $\text{Cl}^-$  transport in lampreys. Nonetheless, it was shown that  $\text{Cl}^-$  transport does support SW-adaptive processes operating within the sea lamprey intestine (Barany et al., 2020). Therefore, it stands unresolved whether an Slc12 family member underlies luminal  $\text{Cl}^-$  removal and supports SW adaptation in this basal vertebrate species (Figure 2).

In contrast to the well-established paradigm for ion secretion by "SW-type" ionocytes, various models seek to explain how "FW-type" ionocytes (within the gill and skin) absorb ions from dilute

environments (Evans, 2008; Takei et al., 2014). These varying models reflect, in part, the evolution of multiple strategies for ion uptake in FW (Dymowska et al., 2012; Takei et al., 2014; Yan & Hwang, 2019). Hiroi et al. (2008) discovered that FW-acclimated Mozambique tilapia express Ncc in the apical membrane of ionocytes. This Ncc is denoted Ncc2 (Slc12a10) and is not a member of the "conventional" Ncc1 (Slc12a3) clade (Takei et al., 2014). Ionocytes that express Ncc2 in tilapia (coined "Type-II" ionocytes) leverage Ncc2 for the transport of  $\text{Na}^+$  and  $\text{Cl}^-$  from FW into the ionocyte interior. Zebrafish (*Danio rerio*) also exhibit Ncc2-mediated  $\text{Cl}^-$  uptake by a subset of ionocytes termed "Ncc-cells" (Wang et al., 2009). Convincing evidence established that Ncc2-expressing ionocytes are in fact critical to  $\text{Cl}^-$  homeostasis (Hiroi et al., 2008; Inokuchi et al., 2008; Kwong & Perry, 2016; Wang et al., 2009). Ncc2-expressing ionocytes operate in species spanning across teleost clades but are seemingly absent in salmonids (Hiroi & McCormick, 2012; Hsu et al., 2014; Inokuchi et al., 2017; Lema et al., 2018). Recently, mummichogs were found to express Ncc2 in the apical region of FW-type ionocytes (Breves et al., 2020). This was a curious finding given that branchial epithelium was previously considered a minor contributor to  $\text{Cl}^-$  uptake by FW-acclimated mummichogs (Marshall et al., 1997; Wood & Laurent, 2003). Rather, the diet may provide  $\text{Cl}^-$  for intestinal absorption with the branchial epithelium operating to minimize  $\text{Cl}^-$  loss via the coordinated expression of tight-junction proteins (Chasiotis et al., 2012; Patrick et al., 1997).

Roles for Ncc in the osmoregulatory organs of basal fishes stand largely undefined. A recent study, however, implicated Ncc in branchial ion absorption by cartilaginous fishes. In Japanese-banded houndshark (*Triakis scyllium*), a "conventional" *ncc1* (*slc12a3*) is expressed in a subpopulation of gill ionocytes (type-B cells) where its expression was increased upon transfer from full SW to 30% SW (Takabe et al., 2016). In turn, Takabe et al. (2016) proposed that for elasmobranchs in which Ncc2 is absent, Ncc1 mediates branchial  $\text{Na}^+$  and  $\text{Cl}^-$  absorption (Figure 2). Similarly, sea lamprey *ncc* is highly expressed in the gill and kidney and is downregulated during acclimation to high-salinity (>50% SW) conditions (Ferreira-Martins et al., 2016). These two studies further indicate that the FW gill Ncc2, which is not expressed in the FW kidney, ought to be regarded as a "ray-finned fish-specific" or "bony fish-specific" Ncc. It follows then that an Ncc2 should be expressed in the gill of FW-acclimated sturgeon. Our preliminary analysis of the sterlet sturgeon genome indicates that two distinct *ncc*-like paralogues exist that have yet to be characterized.

Renal Ncc1 supports  $\text{Na}^+$  and  $\text{Cl}^-$  reabsorption and the production of dilute urine by teleosts inhabiting FW (Kato et al., 2011; Teranishi et al., 2013; Wingert et al., 2007; Figure 2). By localizing Ncc1 to the distal tubules and collecting ducts of FW-acclimated bullshark (*Carcharhinus leucas*), Imaseki et al. (2019) provided the first evidence that Ncc1 contributes to tubular  $\text{Na}^+$  and  $\text{Cl}^-$  reabsorption in elasmobranchs. Finally, hydrochlorothiazide-sensitive transporters such as Ncc1 and Ncc2 were recently proposed as candidate transporters of  $\text{Na}^+$  and  $\text{Cl}^-$  in the desalination of SW by the esophagus of Japanese eel (Takei et al., 2017).

## 2.3 | Anion exchangers (Slc26 family)

Emerging evidence associates Slc26 family anion exchangers with the uptake of environmental  $\text{Cl}^-$  in exchange for  $\text{HCO}_3^-$  by ionocytes (Figure 2). In euryhaline Atlantic stingray (*Dasyatis sabina*), anti-human Slc26a4 (pendrin) immunoreactivity was localized within the branchial epithelium of FW-acclimated animals (Piermarini et al., 2002). In zebrafish, three distinct *slc26a3*, *-4*, and *-6c* genes are expressed by ionocytes. Slc26a4 and *-6c*, however, assume greater roles in  $\text{Cl}^-$  uptake when environmental  $\text{Cl}^-$  is severely depleted (Bayaa et al., 2009; Perry et al., 2009). It remains unclear whether Slc26a3, *-4*, and *-6c* operate within a type of ionocyte that is distinct from the four types already described in zebrafish (HR-, NaR-, Ncc-, or KS-cells; Guh et al., 2015). More recently, it was reported that branchial *slc26a6* expression was higher in FW- versus SW-acclimated rainbow trout (*Oncorhynchus mykiss*), thereby adding additional support for the operation of Slc26-dependent pathways for  $\text{Cl}^-/\text{HCO}_3^-$  exchange in FW-acclimated fishes (Boyle et al., 2015; Leguen et al., 2015). The presence and function of Slc26 family anion exchangers in lampreys and sturgeons remain unresolved and should be the focus of future investigations.

## 2.4 | Voltage-gated $\text{Cl}^-$ channels (Clc $\text{Cl}^-$ channel family members)

Among members of the Clc  $\text{Cl}^-$  channel family, Clc-2, Clc-3, and Clc-K support  $\text{Cl}^-$  balance in fishes (Figure 2). Ion-absorptive ionocytes that express Ncc2 in the apical membrane must transport  $\text{Cl}^-$  across the basolateral membrane. In zebrafish, Clc-2c mediates the movement of  $\text{Cl}^-$  from the interior of Ncc-cells into blood plasma (Pérez-Rius et al., 2015; Wang et al., 2015). Accordingly, knockdown of *clc-2c* perturbed whole-body  $\text{Cl}^-$  balance (Wang et al., 2015). In tilapia, branchial *clc-2c* expression is elevated in parallel with *ncc2* during FW acclimation (Breves et al., 2017). Recall that salmonids lack Ncc2-cells (Hiroi & McCormick, 2012); thus, Clc-2 is not restricted to Ncc2-expressing ionocytes given its gene expression in FW-type rainbow trout ionocytes (Leguen et al., 2015). In pufferfish (*Tetraodon nigroviridis*), European sea bass (*Dicentrarchus labrax*), and Mozambique tilapia, Clc-3 is expressed within branchial ionocytes and may contribute to  $\text{Cl}^-$  uptake from FW (Bossus et al., 2013; Miyazaki et al., 1999; Tang & Lee, 2011; Tang et al., 2010). Within esophageal epithelium, Clc-2 supports the desalination of imbibed SW that enables intestinal water absorption by Japanese eel (Takei et al., 2017). The in vitro effects of serosally applied diphenylamine-2-carboxylic acid (DPC) indicated that Clc-2 mediates the basolateral transport of  $\text{Cl}^-$  by columnar cells within the esophagus (Takei et al., 2017). Finally, Clc-K underlies the capability of renal tubules to reabsorb  $\text{Cl}^-$  in teleosts residing in FW (Kato et al., 2011; Miyazaki et al., 2002; Pérez-Rius et al., 2019; Wingert et al., 2007). Barttin, an accessory subunit that stabilizes Clc-K, is coexpressed with Clc-K in the apical region of distal nephrons of zebrafish (Pérez-Rius et al., 2019). Clc-K is indispensable to  $\text{Cl}^-$  balance given that knockout of *clc-k* is lethal to

embryonic zebrafish undergoing mesonephrogenesis (Pérez-Rius et al., 2019). It currently stands unresolved how zebrafish Clc-K-expressing cells mediate basolateral  $\text{Cl}^-$  transport (Figure 2).

## 3 | ENDOCRINE REGULATION OF $\text{Cl}^-$ TRANSPORT IN FISHES

Perturbations to hydromineral balance elicit the release of hormones that orchestrate the expression, localization, and function of solute transporters/channels. While a broad collection of hormones undoubtedly contributes to  $\text{Cl}^-$  balance in teleosts, prolactin and cortisol have widely accepted roles as “FW- and SW-adapting” hormones, respectively (Takei et al., 2014). We conclude our discussion by highlighting new insights into the roles of prolactin and cortisol (and other corticosteroids) in regulating mechanisms of  $\text{Cl}^-$  transport.

### 3.1 | Prolactin

In agreement with prolactin being recognized as a “FW-adapting” hormone (Pickford & Phillips, 1959), prolactin binding and *prolactin receptor* gene expression occur in key ionoregulatory organs including the gill, kidney, gut, and skin (Breves et al., 2014, 2020). The prolactin receptor was further localized to ionocytes in the gill (Santos et al., 2001; Weng et al., 1997) and to the proximal segment of pronephric tubules (Liu et al., 2006). Only recently have discrete  $\text{Cl}^-$ -transporting processes been linked with prolactin. The failure of hypophysectomized tilapia to adequately recruit Ncc2-expressing ionocytes during FW acclimation was rescued by prolactin replacement (Breves et al., 2010). Prolactin acts directly on the gill and does not require a systemic intermediary to promote Ncc2 expression within ionocytes (Inokuchi et al., 2015). A prolactin-Ncc2 regulatory link also operates in zebrafish and Japanese medaka (*Oryzias latipes*) gill (Bollinger et al., 2018; Breves et al., 2013). In medaka, prolactin stimulated *ncc2* expression via Stat5 activation and not through ERK- or AKT-dependent signal transduction pathways (Bollinger et al., 2018). Moreover, *prolactin*-deficient zebrafish embryos do not fully express *ncc1* and *-2* in the pronephros and epidermis, respectively (Breves et al., 2014; Shu et al., 2016). Given that Ncc2 and Clc-2c are coexpressed within branchial and epidermal ionocytes to facilitate transcellular  $\text{Cl}^-$  transport (Wang et al., 2015), it is logical that prolactin also promotes *clc-2c* expression (Breves et al., 2017; Breves, 2019). To date, any links between prolactin and teleost Slc26s remain entirely uncharacterized.

A distinct *prolactin* gene arose in a vertebrate ancestor that preceded Agnathans (Ocampo Daza & Larhammar, 2018). While only growth hormone has been identified in sea lamprey (Kawauchi et al., 2002), seemingly due to loss of the *prolactin* gene (Ocampo Daza & Larhammar, 2018), sea lamprey express distinct *prolactin* and *growth hormone receptor* genes (Gong et al., 2020). Thus, lampreys are now poised to offer unique insights into how prolactin receptor signaling



(perhaps with growth hormone as the cognate ligand) connects with  $\text{Cl}^-$  balance in a basal vertebrate. Perhaps prolactin receptor-mediated signaling underlies the branchial expression of *ncc* in lampreys residing in FW (Ferreira-Martins et al., 2016).

### 3.2 | Corticosteroids

Cortisol is the primary mineralocorticoid and glucocorticoid in teleosts (Prunet et al., 2006), with both functions mediated primarily via the glucocorticoid receptor. Given these dual roles for cortisol in teleosts, glucocorticoid receptors are widely distributed throughout the body with robust expression in osmoregulatory (gill, kidney, intestine) and gluconeogenic (liver) organs (Greenwood et al., 2003). As an established osmoregulatory hormone in teleosts, cortisol stimulates the abundance and/or activity of ion transporters and pumps that underlie branchial  $\text{Cl}^-$  secretion (e.g., *Nka*, *Nkcc1*, and *Cftr*; McCormick, 2001; Takei et al., 2014). The realization that cortisol can promote branchial ion uptake in some FW-acclimated teleosts arrived long after its SW-adaptive role was firmly established. In the gill and yolk sac of particular teleosts,  $\text{Na}^+/\text{H}^+$  exchanger 3 (*Nhe3*) and *Ncc2* are expressed in distinct ionocytes where they mediate ion absorption (Guh et al., 2015; Hiroi & McCormick, 2012). In larval zebrafish, cortisol stimulated  $\text{Na}^+$  uptake in a fashion dependent upon the presence of *Nhe3b*-expressing cells (Kumai et al., 2012) and promoted the differentiation of *Ncc*-cells from a progenitor population (Cruz et al., 2013; Kumai et al., 2012). Moreover, Lin et al. (2016) showed that cortisol treatment increased *ncc2* expression and *Ncc2*-dependent ion absorption. Cortisol similarly increased *ncc2b* expression in ex vivo Japanese medaka gill (Bossus et al., 2017).

Only recently have roles for corticosteroid signaling towards  $\text{Cl}^-$  balance in basal fishes become clearer. Cortisol is firmly established as a glucocorticoid in sturgeons (Webb et al., 2007). A recent study in Atlantic sturgeon (*Acipenser oxyrinchus oxyrinchus*) reported that plasma cortisol is elevated after SW exposure and promotes branchial *Nka* and *Nkcc1* expression, and, thereby, provided evidence that cortisol is in fact an osmoregulatory hormone in sturgeons (McCormick et al., 2020). Steroidogenic pathways in elasmobranchs result in a novel circulating corticosteroid,  $1\alpha$ -hydroxycorticosterone, which exerts some mineralocorticoid-like activity (Anderson, 2012). Lampreys cannot produce cortisol because they lack  $11\beta$ -hydroxylase; alternatively, 11-deoxycorticosterone (DOC) and the biochemical precursor to cortisol, 11-deoxycortisol (S), are the primary circulating corticosteroids (Close et al., 2010; Rai et al., 2015; Roberts et al., 2014). In vivo administration of S, but not DOC, promoted  $\text{Cl}^-$  homeostasis during a SW challenge by stimulating branchial *Nka* and *Nkcc1*, suggesting that discriminative corticosteroid control of  $\text{Cl}^-$  homeostasis appeared early in vertebrate evolution (Shaughnessy et al., 2020). A functional explanation for the substantial levels of circulating DOC in lamprey is lacking. Although DOC does not appear to control active  $\text{Cl}^-$  transport by the gill (Shaughnessy et al., 2020), it could be that DOC controls tight-junction permeability and barrier

functions in the gill and/or other osmoregulatory organs such as the intestine.

## 4 | CONCLUSION

New perspectives on how epithelial  $\text{Cl}^-$  transport is achieved have fostered the discovery of regulatory connections between the endocrine system and the molecular mechanisms that allow teleost fishes to inhabit diverse aquatic habitats. Prospective studies on basal fishes, which are now enabled by newly available genomic resources, will offer valuable insight into how processes underlying  $\text{Cl}^-$  transport evolved within the vertebrate lineage. As the most basal extant osmoregulating vertebrate, and with its well-developed genome assembly, the sea lamprey will serve as an important model organism in this effort. It will be particularly exciting to unravel how the evolution of vertebrate endocrine systems paralleled the evolution of ionoregulatory strategies.

### ACKNOWLEDGEMENTS

Ciaran A. Shaughnessy is supported by a fellowship provided by the Eugene F. and Easton M. Crawford Charitable Lead Unitrust to National Jewish Health. Jason P. Breves is supported by the National Science Foundation (IOS-1755131).

### CONFLICT OF INTERESTS

The authors declare that there are no conflict of interests.

### DATA AVAILABILITY STATEMENT

Data sharing is not applicable to this article as no new data were created or analyzed in this study.

### ORCID

Ciaran A. Shaughnessy  <http://orcid.org/0000-0003-2146-9126>

Jason P. Breves  <https://orcid.org/0000-0003-1193-4389>

### REFERENCES

- Allen, P. J., Cech, J. J., & Kültz, D. (2009). Mechanisms of seawater acclimation in a primitive, anadromous fish, the green sturgeon. *Journal of Comparative Physiology B*, 179, 903–920.
- Altschul, S. F., Gish, W., Miller, W., Myers, E. W., & Lipman, D. J. (1990). Basic local alignment search tool. *Journal of Molecular Biology*, 215, 403–410.
- Amemiya, C. T., Alföldi, J., Lee, A. P., Fan, S., Philippe, H., MacCallum, I., Braasch, I., Manousaki, T., Schneider, I., Rohner, N., Organ, C., Chalopin, D., Smith, J. J., Robinson, M., Dorrington, R. A., Gerdol, M., Aken, B., Biscotti, M. A., Barucca, M., ... Lindblad-Toh, K. (2013). The African coelacanth genome provides insights into tetrapod evolution. *Nature*, 496, 311–316.
- Anderson, W. G. (2012). The endocrinology of  $1\alpha$ -hydroxycorticosterone in elasmobranch fish: a review. *Comparative Biochemistry and Physiology Part A*, 162, 73–80.
- Ando, M., & Takei, Y. (2015). Guanylin activates  $\text{Cl}^-$  secretion into the lumen of seawater eel intestine via apical  $\text{Cl}^-$  channel under simulated in vivo conditions. *American Journal of Physiology Regulatory, Integrative and Comparative Physiology*, 308, R400–R410.

- Barany, A., Shaughnessy, C. A., Fuentes, J., Mancera, J. M., & McCormick, S. D. (2020). Osmoregulatory role of the intestine in the sea lamprey (*Petromyzon marinus*). *American Journal of Physiology Regulator. Integrative and Comparative Physiology*, 318, R410–R417.
- Bayaa, M., Vulesevic, B., Esbaugh, A., Braun, M., Ekker, M. E., Grosell, M., & Perry, S. F. (2009). The involvement of SLC26 anion transporters in chloride uptake in zebrafish (*Danio rerio*) larvae. *Journal of Experimental Biology*, 212, 3283–3295.
- Bollinger, R. J., Ellis, L. V., Bossus, M. C., & Tipsmark, C. K. (2018). Prolactin controls Na<sup>+</sup>, Cl<sup>-</sup> cotransporter via Stat5 pathway in the teleost gill. *Molecular and Cellular Endocrinology*, 477, 163–171.
- Bossus, M. C., Bollinger, R. J., Reed, P. J., & Tipsmark, C. K. (2017). Prolactin and cortisol regulate branchial claudin expression in Japanese medaka. *General and Comparative Endocrinology*, 240, 77–83.
- Bossus, M., Charmantier, G., Blondeau-Bidet, E., Valletta, B., Boulo, V., & Lorin-Nebel, C. (2013). The CIC-3 chloride channel and osmoregulation in the European sea bass, *Dicentrarchus labrax*. *Journal of Comparative Physiology B*, 183, 641–662.
- Boyle, D., Clifford, A. M., Orr, E., Chamot, D., & Goss, G. G. (2015). Mechanisms of Cl<sup>-</sup> uptake in rainbow trout: cloning and expression of *slc26a6*, a prospective Cl<sup>-</sup>/HCO<sub>3</sub><sup>-</sup> exchanger. *Comparative Biochemistry and Physiology Part A*, 180, 43–50.
- Braasch, I., Gehrke, A. R., Smith, J. J., Kawasaki, K., Manousaki, T., Pasquier, J., Amores, A., Desvignes, T., Batzel, P., Catchen, J., Berlin, A. M., Campbell, M. S., Barrrell, D., Martin, K. J., Mulley, J. F., Ravi, V., Lee, A. P., Nakamura, T., Chalopin, D., ... Postlethwait, J. H. (2016). The spotted gar genome illuminates vertebrate evolution and facilitates human-teleost comparisons. *Nature Genetics*, 48, 427–437.
- Breves, J. P. (2019). Prolactin controls branchial *cln2c* but not *atp1a1a.2* in zebrafish *Danio rerio*. *Journal of Fish Biology*, 94, 168–172.
- Breves, J. P., Keith, P. L. K., Hunt, B. L., Pavlosky, K. K., Inokuchi, M., Yamaguchi, Y., Lerner, D. T., Seale, A. P., & Grau, E. G. (2017). *clc-2c* is regulated by salinity, prolactin and extracellular osmolality in tilapia gill. *Journal of Molecular Endocrinology*, 59, 391–402.
- Breves, J. P., McCormick, S. D., & Karlstrom, R. O. (2014). Prolactin and teleost ionocytes: new insights into cellular and molecular targets of prolactin in vertebrate epithelia. *General and Comparative Endocrinology*, 203, 21–28.
- Breves, J. P., Popp, E. E., Rothenberg, E. F., Rosenstein, C. W., Maffett, K. M., & Guertin, R. R. (2020). Osmoregulatory actions of prolactin in the gastrointestinal tract of fishes. *General and Comparative Endocrinology*, 298, 113589.
- Breves, J. P., Serizier, S. B., Goffin, V., McCormick, S. D., & Karlstrom, R. O. (2013). Prolactin regulates transcription of the ion uptake Na<sup>+</sup>/Cl<sup>-</sup> cotransporter (*ncc*) gene in zebrafish gill. *Molecular and Cellular Endocrinology*, 369, 98–106.
- Breves, J. P., Starling, J. A., Popovski, C. M., Doud, J. M., & Tipsmark, C. K. (2020). Salinity-dependent expression of *ncc2* in opercular epithelium and gill of mummichog (*Fundulus heteroclitus*). *Journal of Comparative Physiology B*, 190, 219–230.
- Breves, J. P., Watanabe, S., Kaneko, T., Hirano, T., & Grau, E. G. (2010). Prolactin restores branchial mitochondrion-rich cells expressing Na<sup>+</sup>/Cl<sup>-</sup> cotransporter in hypophysectomized Mozambique tilapia. *American Journal of Physiology Regulator. Integrative and Comparative Physiology*, 299, R702–R710.
- Chasiotis, H., Kolosov, D., Bui, P., & Kelly, S. P. (2012). Tight junctions, tight junction proteins and paracellular permeability across the gill epithelium of fishes: a review. *Respiratory Physiology and Neurobiology*, 184, 269–281.
- Chen, J. M., Cutler, C., Jacques, C., Boeuf, G., Denamur, E., Lecointre, G., Mercier, B., Cramb, G., & Férec, C. A. (2001). A combined analysis of the cystic fibrosis transmembrane conductance regulator: implications for structure and disease models. *Molecular Biology and Evolution*, 18, 1771–1788.
- Cheng, P., Huang, Y., Du, H., Li, C., Lv, Y., Ruan, R., Ye, H., Bian, C., You, X., Xu, J., Liang, X., Shi, Q., & Wei, Q. (2019). Draft genome and complete *hox*-cluster characterization of the sterlet (*Acipenser ruthenus*). *Frontiers in Genetics*, 10, 1–8.
- Chiu, C. H., Dewar, K., Wagner, G. P., Takahashi, K., Ruddle, F., Ledje, C., Bartsch, P., Scemama, J. L., Stellweg, E., Fried, C., Prohaska, S. J., Stadler, P. F., & Amemiya, C. T. (2004). Bichir HoxA cluster sequence reveals surprising trends in ray-finned fish genomic evolution. *Genome Research*, 14, 11–17.
- Close, D. A., Yun, S. S., McCormick, S. D., Wildbill, A. J., & Li, W. (2010). 11-deoxycortisol is a corticosteroid hormone in the lamprey. *Proceedings of the National Academy of Sciences United States of America*, 107, 13942–13947.
- Cruz, S. A., Chao, P. L., & Hwang, P. P. (2013). Cortisol promotes differentiation of epidermal ionocytes through Foxi3 transcription factors in zebrafish (*Danio rerio*). *Comparative Biochemistry and Physiology Part A*, 164, 249–257.
- Cui, G., Hong, J., Chung-Davidson, Y. W., Infield, D., Xu, X., Li, J., Simhaev, L., Khazanov, N., Stauffer, B., Imhoff, B., Cottrill, K., Blalock, J. E., Li, W., Senderowitz, H., Sorscher, E., McCarty, N. A., & Gaggari, A. (2019). An ancient CFTR ortholog informs molecular evolution in ABC transporters. *Developmental Cell*, 51, 421–430.
- Du, K., Stöck, M., Kneitz, S., Klopp, C., Woltering, J. M., Adolphi, M. C., Feron, R., Prokopov, D., Makunin, A., Kichigin, I., Schmidt, C., Fischer, P., Kuhl, H., Wuertz, S., Gessner, J., Kloas, W., Cabau, C., lampietro, C., Parrinello, H., ... Scharf, M. (2020). The sterlet sturgeon genome sequence and the mechanisms of segmental rediploidization. *Nature Ecology & Evolution*, 4, 841–852.
- Dymowska, A. K., Hwang, P. P., & Goss, G. G. (2012). Structure and function of ionocytes in the freshwater fish gill. *Respiratory Physiology and Neurobiology*, 184, 282–292.
- Evans, D. H. (2008). Teleost fish osmoregulation: what have we learned since August Krogh, Homer Smith, and Ancel Keys. *American Journal of Physiology: Regulatory, Integrative and Comparative Physiology*, 295, R704–R713.
- Ferreira-Martins, D., Coimbra, J., Antunes, C., & Wilson, J. M. (2016). Effects of salinity on upstream-migrating, spawning sea lamprey. *Petromyzon marinus. Conservation Physiology*, 4, cov064.
- Gagnon, E., Forbush, B., Flemmer, A. W., Giménez, I., Caron, L., & Isenring, P. (2002). Functional and molecular characterization of the shark renal Na-K-Cl cotransporter: novel aspects. *American Journal of Physiology. Renal Physiology*, 283, F1046–F1055.
- Gong, N., Ferreira-Martins, D., McCormick, S. D., & Sheridan, M. A. (2020). Divergent genes encoding the putative receptors for growth hormone and prolactin in sea lamprey display distinct patterns of expression. *Scientific Reports*, 10, 1674.
- Greenwood, A. K., Butler, P. C., White, R. B., Demarco, U., Pearce, D., & Fernald, R. D. (2003). Multiple corticosteroid receptors in a teleost fish: Distinct sequences, expression patterns, and transcriptional activities. *Endocrinology*, 144, 4226–4236.
- Gregório, S. F., Carvalho, E. S., Encarnação, S., Wilson, J. M., Power, D. M., Canário, A. V., & Fuentes, J. (2013). Adaptation to different salinities exposes functional specialization in the intestine of the sea bream (*Sparus aurata* L.). *Journal of Experimental Biology*, 216, 470–479.
- Grosell, M. (2006). Intestinal anion exchange in marine fish osmoregulation. *Journal of Experimental Biology*, 209, 2813–2827.
- Grosell, M. (2014). The physiology of fishes. In D. H. Evans, J. B. Claiborne, & S. Currie (Eds.), *Intestinal transport* (4th Ed., pp. 175–203). Boca Raton, FL: CRC Press.
- Guh, Y. J., Lin, C. H., & Hwang, P. P. (2015). Osmoregulation in zebrafish: ion transport mechanisms and functional regulation. *EXCLI Journal*, 14, 627–659.
- Hiroi, J., & McCormick, S. D. (2012). New insights into gill ionocyte and ion transporter function in euryhaline and diadromous fish. *Respiratory Physiology and Neurobiology*, 184, 257–268.

- Hiroi, J., Yasumasu, S., McCormick, S. D., Hwang, P. P., & Kaneko, T. (2008). Evidence for an apical Na-Cl cotransporter involved in ion uptake in a teleost fish. *Journal of Experimental Biology*, 211, 2584–2599.
- Hsu, H. H., Lin, L. Y., Tseng, Y. C., Horng, J. L., & Hwang, P. P. (2014). A new model for fish ion regulation: identification of ionocytes in freshwater- and seawater-acclimated medaka (*Oryzias latipes*). *Cell and Tissue Research*, 357, 225–243.
- Imaseki, I., Wakabayashi, M., Hara, Y., Watanabe, T., Takabe, S., Kakumura, K., Honda, Y., Ueda, K., Murakumo, K., Matsumoto, R., Matsumoto, Y., Nakamura, M., Takagi, W., Kuraku, S., & Hyodo, S. (2019). Comprehensive analysis of genes contributing to euryhalinity in the bull shark, *Carcharhinus leucas*; Na<sup>+</sup>-Cl<sup>-</sup> co-transporter is one of the key renal factors upregulated in acclimation to low-salinity environment. *Journal of Experimental Biology*, 222, jeb201780.
- Inokuchi, M., Breves, J. P., Moriyama, S., Watanabe, S., Kaneko, T., Lerner, D. T., Grau, E. G., & Seale, A. P. (2015). Prolactin 177, prolactin 188, and extracellular osmolality independently regulate the gene expression of ion transport effectors in gill of Mozambique tilapia. *American Journal of Physiology: Regulatory, Integrative and Comparative Physiology*, 309, R1251–R1263.
- Inokuchi, M., Hiroi, J., Watanabe, S., Lee, K. M., & Kaneko, T. (2008). Gene expression and morphological localization of NHE3, NCC and NKCC1a in branchial mitochondria-rich cells of Mozambique tilapia (*Oreochromis mossambicus*) acclimated to a wide range of salinities. *Comparative Biochemistry and Physiology Part A*, 151, 151–158.
- Inokuchi, M., Nakamura, M., Miyanishi, H., Hiroi, J., & Kaneko, T. (2017). Functional classification of gill ionocytes and spatiotemporal changes in their distribution after transfer from seawater to freshwater in Japanese seabass. *Journal of Experimental Biology*, 220, 4720–4732.
- Karnaky, K. J., Degnan, K. J., & Zadunaisky, J. A. (1977). Chloride transport across isolated opercular epithelium of killifish: a membrane rich in chloride cells. *Science*, 195(4274), 203–205.
- Kato, A., Muro, T., Kimura, Y., Li, S., Islam, Z., Ogoshi, M., Doi, H., & Hirose, S. (2011). Differential expression of Na<sup>+</sup>-Cl<sup>-</sup> cotransporter and Na<sup>+</sup>-K<sup>+</sup>-Cl<sup>-</sup> cotransporter 2 in the distal nephrons of euryhaline and seawater pufferfishes. *American Journal of Physiology Regulatory, Integrative and Comparative Physiology*, 300, R284–R297.
- Kawauchi, H., Suzuki, K., Yamazaki, T., Moriyama, S., Nozaki, M., Yamaguchi, K., Takahashi, A., Youson, J., & Sower, S. A. (2002). Identification of growth hormone in the sea lamprey, an extant representative of a group of the most ancient vertebrates. *Endocrinology*, 143, 4916–4921.
- Kumai, Y., Nesan, D., Vijayan, M. M., & Perry, S. F. (2012). Cortisol regulates Na<sup>+</sup> uptake in zebrafish, *Danio rerio*, larvae via the glucocorticoid receptor. *Molecular and Cellular Endocrinology*, 364, 113–125.
- Kwong, R. W., & Perry, S. F. (2016). A role for sodium-chloride cotransporters in the rapid regulation of ion uptake following acute environmental acidosis: new insights from the zebrafish model. *American Journal of Physiology Cell Physiology*, 311, C931–C941.
- Leguen, I., Le Cam, A., Montfort, J., Peron, S., & Fautrel, A. (2015). Transcriptomic analysis of trout gill ionocytes in fresh water and sea water using laser capture microdissection combined with microarray analysis. *PLOS One*, 10, e0139938.
- Lema, S. C., Carvalho, P. G., Egelston, J. N., Kelly, J. T., & McCormick, S. D. (2018). Dynamics of gene expression responses for ion transport proteins and aquaporins in the gill of a euryhaline pupfish during freshwater and high-salinity acclimation. *Physiological and Biochemical Zoology*, 91, 1148–1171.
- Li, Z., Lui, E. Y., Wilson, J. M., Ip, Y. K., Lin, Q., Lam, T. J., & Lam, S. H. (2014). Expression of key ion transporters in the gill and esophageal-gastrointestinal tract of euryhaline Mozambique tilapia *Oreochromis mossambicus* acclimated to fresh water, seawater and hypersaline water. *PLOS One*, 9, e87591.
- Lin, C. H., Hu, H. J., & Hwang, P. P. (2016). Cortisol regulates sodium homeostasis by stimulating the transcription of sodium-chloride transporter (NCC) in zebrafish (*Danio rerio*). *Molecular and Cellular Endocrinology*, 422, 93–102.
- Liu, N. A., Liu, Q., Wawrowsky, K., Yang, Z., Lin, S., & Melmed, S. (2006). Prolactin receptor signaling mediates the osmotic response of embryonic zebrafish lactotrophs. *Molecular Endocrinology*, 20, 871–880.
- Marshall, W. S., & Bellamy, D. (2010). The 50 year evolution of in vitro systems to reveal salt transport functions of teleost fish gills. *Comparative Biochemistry and Physiology Part A*, 155, 275–280.
- Marshall, W. S., Bryson, S. E., Darling, P., Whitten, C., Patrick, M., Wilkie, M., Wood, C. M., & Buckland Nicks, J. (1997). NaCl transport and ultrastructure of opercular epithelium from a freshwater-adapted euryhaline teleost, *Fundulus heteroclitus*. *Journal of Experimental Zoology*, 277, 23–37.
- Marshall, W. S., & Grosell, M. (2006). The physiology of fishes. In D. H. Evans, & J. B. Claiborne (Eds.), *Ion transport, osmoregulation, and acid-base balance* (3rd Ed., pp. 177–230). Boca Raton, FL: CRC Press.
- Marshall, W. S., Howard, J. A., Cozzi, R. R., & Lynch, E. M. (2002). NaCl and fluid secretion by the intestine of the teleost *Fundulus heteroclitus*: involvement of CFTR. *Journal of Experimental Biology*, 205, 745–758.
- Marshall, W. S., & Singer, T. D. (2002). Cystic fibrosis transmembrane conductance regulator in teleost fish. *Biochimica et Biophysica Acta*, 1566, 16–27.
- McCormick, S. D. (2001). Endocrine control of osmoregulation in teleost fish. *American Zoologist*, 41, 781–794.
- McCormick, S. D., Taylor, M. L., & Regish, A. M. (2020). Cortisol is an osmoregulatory and glucose-regulating hormone in Atlantic sturgeon, a basal ray-finned fish. *Journal of Experimental Biology*, 223, jeb220251.
- Miyazaki, H., Kaneko, T., Uchida, S., Sasaki, S., & Takei, Y. (2002). Kidney-specific chloride channel, OmCIC-K, predominantly expressed in the diluting segment of freshwater-adapted tilapia kidney. *Proceedings of the National Academy of Sciences United States of America*, 99, 15782–15787.
- Miyazaki, H., Uchida, S., Takei, Y., Hirano, T., Marumo, F., & Sasaki, S. (1999). Molecular cloning of CLC chloride channels in *Oreochromis mossambicus* and their functional complementation of yeast CLC gene mutant. *Biochemical and Biophysical Research Communications*, 255, 175–181.
- Ocampo Daza, D., & Larhammar, D. (2018). Evolution of the growth hormone, prolactin, prolactin 2 and somatolactin family. *General and Comparative Endocrinology*, 264, 94–112.
- Patrick, M. L., Pärt, P., Marshall, W. S., & Wood, C. M. (1997). Characterization of ion and acid-base transport in the freshwater adapted mummichog (*Fundulus heteroclitus*). *Journal of Experimental Zoology*, 279, 208–219.
- Pelis, R. M., Zydlewski, J., & McCormick, S. D. (2001). Gill Na<sup>+</sup>-K<sup>+</sup>-2Cl<sup>-</sup> cotransporter abundance and location in Atlantic salmon: effects of seawater and smolting. *American Journal of Physiology Regulatory, Integrative and Comparative Physiology*, 280, R1844–R1852.
- Pérez-Rius, C., Castellanos, A., Gaitán-Peñas, H., Navarro, A., Artuch, R., Barrallo-Gimeno, A., & Estévez, R. (2019). Role of zebrafish CIC-K/ barttin channels in apical kidney chloride reabsorption. *Journal of Physiology*, 597, 3969–3983.
- Pérez-Rius, C., Gaitán-Peñas, H., Estévez, R., & Barrallo-Gimeno, A. (2015). Identification and characterization of the zebrafish CIC-2 chloride channel orthologs. *Pflügers Archiv*, 467, 1769–1781.
- Perry, S. F., Vulesevic, B., Grosell, M., & Bayaa, M. (2009). Evidence that SLC26 anion transporters mediate branchial chloride uptake in adult zebrafish (*Danio rerio*). *American Journal of Physiology Regulatory, Integrative and Comparative Physiology*, 297, R988–R997.
- Pickford, G. E., & Phillips, J. G. (1959). Prolactin, a factor in promoting survival of hypophysectomized killifish in fresh water. *Science*, 130, 454–455.



- Piermarini, P. M., Verlander, J. W., Royaux, I. E., & Evans, D. H. (2002). Pendrin immunoreactivity in the gill epithelium of a euryhaline elasmobranch. *American Journal of Physiology Regulatory, Integrative and Comparative Physiology*, 283, R983–R992.
- Prunet, P., Sturm, A., & Milla, S. (2006). Multiple corticosteroid receptors in fish: from old ideas to new concepts. *General and Comparative Endocrinology*, 147, 17–23.
- Rai, S., Szeitz, A., Roberts, B. W., Christie, Q., Didier, W., Eom, J., Yun, S. S., & Close, D. A. (2015). A putative corticosteroid hormone in Pacific lamprey, *Entosphenus tridentatus*. *General and Comparative Endocrinology*, 212, 178–184.
- Ren, J., Chung-Davidson, Y. W., Yeh, C. Y., Scott, C., Brown, T., & Li, W. (2015). Genome-wide analysis of the ATP-binding cassette (ABC) transporter gene family in sea lamprey and Japanese lamprey. *BMC Genomics*, 16, 436.
- Roberts, B. W., Didier, W., Rai, S., Johnson, N. S., Libants, S., Yun, S. S., & Close, D. A. (2014). Regulation of a putative corticosteroid, 17,21-dihydroxypregn-4-ene,3,20-one, in sea lamprey, *Petromyzon marinus*. *General and Comparative Endocrinology*, 196, 17–25.
- Santos, C. R. A., Ingleton, P. M., Cavaco, J. E. B., Kelly, P. A., Edery, M., & Power, D. M. (2001). Cloning, characterization, and tissue distribution of prolactin receptor in the sea bream (*Sparus aurata*). *General and Comparative Endocrinology*, 121, 32–47.
- Sardella, B. A., & Kültz, D. (2009). Osmo- and ionoregulatory responses of green sturgeon (*Acipenser medirostris*) to salinity acclimation. *Journal of Comparative Physiology B*, 179, 383–390.
- Shaughnessy, C. A., Baker, D. W., Brauner, C. J., Morgan, J. D., & Bystriansky, J. S. (2015). Interaction of osmoregulatory and acid-base compensation in white sturgeon (*Acipenser transmontanus*) during exposure to aquatic hypercarbia and elevated salinity. *Journal of Experimental Biology*, 218, 2712–2719.
- Shaughnessy, C. A., Barany, A., & McCormick, S. D. (2020). 11-Deoxycortisol controls hydromineral balance in the most basal osmoregulating vertebrate, sea lamprey (*Petromyzon marinus*). *Scientific Reports*, 10, 12148.
- Shaughnessy, C. A., & McCormick, S. D. (2020). Functional characterization and osmoregulatory role of the Na<sup>+</sup>-K<sup>+</sup>-2Cl<sup>-</sup> cotransporter in the gill of sea lamprey (*Petromyzon marinus*), a basal vertebrate. *American Journal of Physiology: Regulatory, Integrative and Comparative Physiology*, 318, R17–R29.
- Shu, Y., Lou, Q., Dai, Z., Dai, X., He, J., Hu, W., & Yin, Z. (2016). The basal function of teleost prolactin as a key regulator on ion uptake identified with zebrafish knockout models. *Scientific Reports*, 6, 18597.
- Singer, T. D., Tucker, S. J., Marshall, W. S., & Higgins, C. F. (1998). A divergent CFTR homologue: highly regulated salt transport in the euryhaline teleost *F. heteroclitus*. *American Journal of Physiology Cell Physiology*, 274, C715–C723.
- Smith, J. J., Kuraku, S., Holt, C., Sauka-Spengler, T., Jiang, N., Campbell, M. S., Yandell, M. D., Manousaki, T., Meyer, A., Bloom, O. E., Morgan, J. R., Buxbaum, J. D., Sachidanandam, R., Sims, C., Garruss, A. S., Cook, M., Krumlauf, R., Wiedemann, L. M., Sower, S. A., ... Li, W. (2013). Sequencing of the sea lamprey (*Petromyzon marinus*) genome provides insights into vertebrate evolution. *Nature Genetics*, 45, 415–421.
- Smith, J. J., Timoshevskaya, N., Ye, C., Holt, C., Keinath, M. C., Parker, H. J., Cook, M. E., Hess, J. E., Narum, S. R., Lamanna, F., Kaessmann, H., Timoshevskiy, V. A., Waterbury, C. K. M., Saraceno, C., Wiedemann, L. M., Robb, S. M. C., Baker, C., Eichler, E. E., Hockman, D., ... Amemiya, C. T. (2018). The sea lamprey germline genome provides insights into programmed genome rearrangement and vertebrate evolution. *Nature Genetics*, 50, 270–277.
- Sundell, K., & Sundh, H. (2012). Intestinal fluid absorption in anadromous salmonids: importance of tight junctions and aquaporins. *Frontiers in Physiology*, 3, 338.
- Sundh, H., Nilsen, T. O., Lindström, J., Hasselberg-Frank, L., Stefansson, S. O., McCormick, S. D., & Sundell, K. (2014). Development of intestinal ion-transporting mechanisms during smoltification and seawater acclimation in Atlantic salmon *Salmo salar*. *Journal of Fish Biology*, 85, 1227–1252.
- Takabe, S., Inokuchi, M., Yamaguchi, Y., & Hyodo, S. (2016). Distribution and dynamics of branchial ionocytes in houndshark reared in full-strength and diluted seawater environments. *Comparative Biochemistry and Physiology Part A*, 198, 22–32.
- Takei, Y., Hiroi, J., Takahashi, H., & Sakamoto, T. (2014). Diverse mechanisms for body fluid regulation in teleost fishes. *American Journal of Physiology Regulatory, Integrative and Comparative Physiology*, 307, R778–R792.
- Takei, Y., Wong, M. K., Pipil, S., Ozaki, H., Suzuki, Y., Iwasaki, W., & Kusakabe, M. (2017). Molecular mechanisms underlying active desalination and low water permeability in the esophagus of eels acclimated to seawater. *American Journal of Physiology Regulatory, Integrative and Comparative Physiology*, 312, R231–R244.
- Tang, C. H., Hwang, L. Y., & Lee, T. H. (2010). Chloride channel CIC-3 in gills of the euryhaline teleost, *Tetraodon nigroviridis*: expression, localization and the possible role of chloride absorption. *Journal of Experimental Biology*, 213, 683–693.
- Tang, C. H., & Lee, T. H. (2011). Ion-deficient environment induces the expression of basolateral chloride channel, CIC-3-like protein, in gill mitochondrion-rich cells for chloride uptake of the tilapia *Oreochromis mossambicus*. *Physiological and Biochemical Zoology*, 84, 54–67.
- Teranishi, K., Mekuchi, M., & Kaneko, T. (2013). Expression of sodium/hydrogen exchanger 3 and cation-chloride cotransporters in the kidney of Japanese eel acclimated to a wide range of salinities. *Comparative Biochemistry and Physiology Part A*, 164, 333–343.
- Vialle, R. A., De Souza, J. E. S., De Paiva Lopes, K., Teixeira, D. G., De Azevedo Alves Sobrinho, P., Ribeiro-Dos-Santos, A. M., Furtado, C., Sakamoto, T., Silva, F. A. O., De Oliveira, E. H. C., Hamoy, I. G., Assumpção, P. P., Ribeiro-Dos-Santos, Á., Lima, J. P. M. S., Seuanez, H. N., De Souza, S. J., & Santos, S. (2018). Whole genome sequencing of the pirarucu (*Arapaima gigas*) supports independent emergence of major teleost clades. *Genome Biology and Evolution*, 10, 2366–2379.
- Wang, Y. F., Tseng, Y. C., Yan, J. J., Hiroi, J., & Hwang, P. P. (2009). Role of SLC12A10.2, a Na-Cl cotransporter-like protein, in a Cl uptake mechanism in zebrafish (*Danio rerio*). *American Journal of Physiology: Regulatory, Integrative and Comparative Physiology*, 296, R1650–R1660.
- Wang, Y. F., Yan, J. J., Tseng, Y. C., Chen, R. D., & Hwang, P. P. (2015). Molecular physiology of an extra-renal Cl<sup>-</sup> uptake mechanism for body fluid Cl<sup>-</sup> homeostasis. *International Journal of Biological Sciences*, 11, 1190–1203.
- Webb, M. A., Allert, J. A., Kappenman, K. M., Marcos, J., Feist, G. W., Schreck, C. B., & Shackleton, C. H. (2007). Identification of plasma glucocorticoids in pallid sturgeon in response to stress. *General and Comparative Endocrinology*, 154, 98–104.
- Weng, C. F., Lee, T. H., & Hwang, P. P. (1997). Immune localization of prolactin receptor the mitochondria-rich cells of the euryhaline teleost (*Oreochromis mossambicus*) gill. *FEBS Letters*, 405, 91–94.
- Whittamore, J. M. (2012). Osmoregulation and epithelial water transport: lessons from the intestine of marine teleost fish. *Journal of Comparative Physiology B*, 182, 1–39.
- Wingert, R. A., Selleck, R., Yu, J., Song, H. D., Chen, Z., Song, A., Zhou, Y., Thisse, B., Thisse, C., McMahon, A. P., & Davidson, A. J. (2007). The *cdx* genes and retinoic acid control the positioning and segmentation of the zebrafish pronephros. *PLOS Genetics*, 3, 1922–1938.
- Wong, M. K., Pipil, S., Kato, A., & Takei, Y. (2016). Duplicated CFTR isoforms in eels diverged in regulatory structures and osmoregulatory functions. *Comparative Biochemistry and Physiology Part A*, 199, 130–141.
- Wood, C. M., & Laurent, P. (2003). Na<sup>+</sup> versus Cl<sup>-</sup> transport in the intact killifish after rapid salinity transfer. *Biochimica et Biophysica Acta*, 1618, 106–119.

- Yamaguchi, K., Hara, Y., Tatsumi, K., Nishimura, O., Smith, J. J., Kadota, M., & Kuraku, S. (2020). Inference of a genome-wide protein-coding gene set of the inshore hagfish *Eptatretus burgeri*. *bioRxiv*. <https://www.biorxiv.org/content/10.1101/2020.07.24.218818v1>
- Yan, J. J., & Hwang, P. P. (2019). Novel discoveries in acid-base regulation and osmoregulation: a review of selected hormonal actions in zebrafish and medaka. *General and Comparative Endocrinology*, 277, 20–29.

**How to cite this article:** Shaughnessy CA, Breves JP. Molecular mechanisms of Cl<sup>-</sup> transport in fishes: New insights and their evolutionary context. *J. Exp. Zool.* 2021; 335:207–216. <https://doi.org/10.1002/jez.2428>